


Farmland Soil Map

2-E-24-SP / 2-J-24-SP - 6701 Roberts Rd

65.3% is prime or locally important farmland soil - 48.39 acres



KGIS makes no representation or warranty as to the accuracy of this map and its information, nor to its fitness for use. Any user of this map product accepts the same AS IS, WITH ALL FAULTS, and assumes all responsibility for the use thereof, and further covenants and agrees to hold KGIS harmless from any and all damage, loss, or liability arising from any use of this map product.

© KGIS 2024

Printed: 4/19/2024 6:28:07 PM

Prime and Other Important Farmlands

Knox County, Tennessee

Map symbol	Map unit name	Farmland classification
CoB	Corryton loam, 2 to 5 percent slopes	All areas are prime farmland
CtB	Corryton-Townley complex, 2 to 5 percent slopes	All areas are prime farmland
DeB	Dewey silt loam, 2 to 6 percent slopes	All areas are prime farmland
EmB	Emory silt loam, 2 to 5 percent slopes	All areas are prime farmland
EvB	Etowah-Minvale complex, 2 to 5 percent slopes	All areas are prime farmland
He	Heiskell silt loam, 0 to 2 percent slopes, occasionally flooded	All areas are prime farmland
HeB	Heiskell silt loam, 2 to 5 percent slopes	All areas are prime farmland
Ph	Pettyjon-Hamblen complex, 0 to 3 percent slopes, occasionally flooded	All areas are prime farmland
ShB	Shady loam, 2 to 5 percent slopes	All areas are prime farmland
So	Shady-Whitwell complex, 0 to 3 percent slopes, rarely flooded	All areas are prime farmland
St	Steadman silt loam, 0 to 3 percent slopes, occasionally flooded	All areas are prime farmland
SwB	Swafford silt loam, 2 to 5 percent slopes	All areas are prime farmland
Tc	Toccoa fine sandy loam, 0 to 3 percent slopes, rarely flooded	All areas are prime farmland
WeB	Waynesboro loam, 2 to 6 percent slopes	All areas are prime farmland
WwB	Whitwell loam, 2 to 5 percent slopes, rarely flooded	All areas are prime farmland
CcC	Coghill-Corryton complex, 5 to 12 percent slopes	Farmland of local importance
CkC	Collegedale-Talbott complex, 5 to 12 percent slopes, rocky	Farmland of local importance
CoC	Corryton loam, 5 to 12 percent slopes	Farmland of local importance
CtC	Corryton-Townley complex, 5 to 12 percent slopes	Farmland of local importance
DeC2	Dewey silt loam, 6 to 15 percent slopes, eroded	Farmland of local importance
EmC	Emory silt loam, 5 to 12 percent slopes	Farmland of local importance
FvC	Fullerton-Minvale complex, 5 to 12 percent slopes	Farmland of local importance
ShC	Shady loam, 5 to 12 percent slopes	Farmland of local importance
SwC	Swafford silt loam, 5 to 12 percent slopes	Farmland of local importance
ToC	Townley silt loam, 5 to 12 percent slopes	Farmland of local importance
WeC	Waynesboro loam, 6 to 15 percent slopes	Farmland of local importance

Prime and Other Important Farmlands

This table lists the map units in the survey area that are considered important farmlands. Important farmlands consist of prime farmland, unique farmland, and farmland of statewide or local importance. This list does not constitute a recommendation for a particular land use.

In an effort to identify the extent and location of important farmlands, the Natural Resources Conservation Service, in cooperation with other interested Federal, State, and local government organizations, has inventoried land that can be used for the production of the Nation's food supply.

"Prime farmland" is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil quality, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. The water supply is dependable and of adequate quality. Prime farmland is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

For some of the soils identified in the table as prime farmland, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures.

A recent trend in land use in some areas has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

"Unique farmland" is land other than prime farmland that is used for the production of specific high-value food and fiber crops, such as citrus, tree nuts, olives, cranberries, and other fruits and vegetables. It has the special combination of soil quality, growing season, moisture supply, temperature, humidity, air drainage, elevation, and aspect needed for the soil to economically produce sustainable high yields of these crops when properly managed. The water supply is dependable and of adequate quality. Nearness to markets is an additional consideration. Unique farmland is not based on national criteria. It commonly is in areas where there is a special microclimate, such as the wine country in California.

In some areas, land that does not meet the criteria for prime or unique farmland is considered to be "farmland of statewide importance" for the production of food, feed, fiber, forage, and oilseed crops. The criteria for defining and delineating farmland of statewide importance are determined by the appropriate State agencies. Generally, this land includes areas of soils that nearly meet the requirements for prime farmland and that economically produce high yields of crops when treated and managed according to acceptable farming methods. Some areas may produce as high a yield as prime farmland if conditions are favorable. Farmland of statewide importance may include tracts of land that have been designated for agriculture by State law.

In some areas that are not identified as having national or statewide importance, land is considered to be "farmland of local importance" for the production of food, feed, fiber, forage, and oilseed crops. This farmland is identified by the appropriate local agencies. Farmland of local importance may include tracts of land that have been designated for agriculture by local ordinance.